



## VACCINES REQUIRED FOR SCHOOL ATTENDANCE, GRADES K-12

July 1, 2017 – June 30, 2018

VACCINE	Kindergarten - 6 <sup>th</sup> Grade	7 <sup>th</sup> - 10 <sup>th</sup> Grade	11 <sup>th</sup> - 12 <sup>th</sup> Grade
<b>Hepatitis B</b>	<b>3 doses</b> Dose 3 must be given on or after 24 weeks of age		Dose 3 must be given on or after 4 months of age
<b>Diphtheria, Tetanus, and Pertussis (DTaP/DT/Td/Tdap)</b>	<b>5 doses</b> (4 doses only <b>IF</b> 4 <sup>th</sup> dose given on or after 4 <sup>th</sup> birthday)  <i>Plus</i>  <b>1 dose Tdap required for 6<sup>th</sup>-12<sup>th</sup> grade AND</b> on or after 11 years of age <i>(see page 2 for more details)</i>		
<b>Polio</b> (IPV or OPV)	<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given on or after 4 <sup>th</sup> birthday) <ul style="list-style-type: none"> <li>• The final dose given on or after August 7, 2009, must be given on or after 4 years of age AND a minimum interval of 6 months from the previous dose.</li> </ul>	<b>4 doses</b> (3 doses only <b>IF</b> 3 <sup>rd</sup> dose given on or after 4 <sup>th</sup> birthday)	
<b>Measles, Mumps, and Rubella</b>	<b>2 doses</b>		
<b>Varicella</b>	<b>2 doses</b>  <b>OR</b>  Healthcare provider verified disease		

- Look at the Minimum Age and Interval Table on page 2 for recommended minimum age and spacing information.
- Review the Individual Vaccine Requirements Summary for more detailed information:  
[www.doh.wa.gov/CommunityandEnvironment/Schools/Immunization/VaccineRequirements.aspx](http://www.doh.wa.gov/CommunityandEnvironment/Schools/Immunization/VaccineRequirements.aspx)

## Minimum Age & Interval for Valid Vaccine Doses

Vaccine	Dose #	Minimum Age	Minimum Interval Between Doses	Notes
Hepatitis B <b>HepB</b>	Dose 1	Birth	4 weeks between Dose 1 & 2 (K-12 <sup>th</sup> )	<ul style="list-style-type: none"> <li>2 doses valid if adult Recombivax HB® given between ages 11 and 15 and doses separated by at least 4 months.</li> </ul>
	Dose 2	4 weeks	8 weeks between Dose 2 & 3 (K-12 <sup>th</sup> )	
	Dose 3	24 weeks 4 months	16 weeks between Dose 1 & 3 (K-10 <sup>th</sup> ) 12 weeks between Dose 1 & 3 (11 <sup>th</sup> -12 <sup>th</sup> )	
Diphtheria, Tetanus, and Pertussis <b>DTaP/DT</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>DTaP: for children through age 6.</li> <li>6 month interval is recommended between Dose 3 and Dose 4, but minimum interval of 4 months is acceptable.</li> <li>Students 7-10 years of age not fully immunized with DTaP should get one Tdap followed by additional doses of Td if needed.</li> <li>DTaP given after age 7 counts for the Tdap dose; no Tdap required at 11-12 years of age.</li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	12 months	6 months between Dose 4 & 5	
	Dose 5	4 years	–	
Tetanus, Diphtheria, and Pertussis <b>Tdap</b>	Dose 1	10 years recommended. See notes for exceptions	–	<ul style="list-style-type: none"> <li>Tdap: for children 7 years of age or older.</li> <li>If no DTaP doses given before age 7, students must get Tdap followed by 2 doses of Td.</li> <li>Tdap given between 7-10 years of age is valid and meets the requirement.</li> <li>Can be given regardless of the interval between DTaP or Td.</li> </ul>
Tetanus and Diphtheria <b>Td</b>	Dose 1	7 years	5 years	<ul style="list-style-type: none"> <li>Td: for children 7 years of age or older.</li> </ul>
Polio <b>IPV or OPV</b>	Dose 1	6 weeks	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>Not required for students 18 years and older.</li> <li>If 4 doses of OPV received, and all doses given before 4 years of age, one dose of IPV is required at 4 years and older.</li> <li>Please see Individual Vaccine Requirements Summary for more details: <a href="http://www.doh.wa.gov/CommunityandEnvironment/Schools/Immunization/VaccineRequirements.aspx">www.doh.wa.gov/CommunityandEnvironment/Schools/Immunization/VaccineRequirements.aspx</a></li> </ul>
	Dose 2	10 weeks	4 weeks between Dose 2 & 3	
	Dose 3	14 weeks	6 months between Dose 3 & 4	
	Dose 4	4 years	–	
Measles, Mumps, and Rubella <b>MMR</b>	Dose 1	12 months	4 weeks between Dose 1 & 2	<ul style="list-style-type: none"> <li>MMRV (MMR + varicella) may be used instead of separate MMR and varicella vaccines.</li> <li>Must get the same day as VAR <b>OR</b> at least 28 days apart.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine such as MMR and MMR. The 4 day grace period <b>DOES NOT</b> apply between Dose 1 and Dose 2 of different live vaccines, such as between MMR and Varicella or between MMR and live flu vaccine.</li> </ul>
	Dose 2	13 months	–	
Varicella (chickenpox) <b>VAR</b>	Dose 1	12 months	3 months between Dose 1 & 2 (12 months through 12 years) 4 weeks between Dose 1 & 2 (13 years and older)	<ul style="list-style-type: none"> <li>Recommended: 3 months between varicella doses, but minimum interval of 28 days acceptable. Minimum age of 13 months also acceptable.</li> <li>Must get the same day as MMR <b>OR</b> at least 28 days apart.</li> <li>4-day grace <b>DOES</b> apply between doses of the same live vaccine; <b>DOES NOT</b> apply between doses of different live vaccines.</li> </ul>
	Dose 2	15 months	–	